Author Index

Abe, Y.
——, Kobayashi, E., Suzuki, N., Kaneko, A., Fujie, T., Nagata, Y., Matsushima, Y., Takai, N., Watanabe, T. and Otsubo, O.
Multi-component analysis of urine of dogs with liver diseases by high-performance liquid chromatography 11
Adamcová, E., see Kříž, J. 159
Alvaro, D., see Cantafora, A. 367
Angelico, M., see Cantafora, A. 367
Báti, J., see Kató, K. 345
Belle, H. van, see Wynants, J. 297
Biggs, W. R., see Fetzer, J. C. 87
——, see Jinno, K. 123
Bosch, I., see Brückner, H. 251
Brückner, H.
——, Bosch, I., Graser, T. and Fürst, P.
Determination of non-protein α-alkyl-α-amino acids by reversed-phase high-performance liquid chromatography in comparison with ion-exchange and capillary gas chromatography 251
Burda, J., see Kříž, J. 159
Calabrese, K. L.
Analysis of fatty acids in lubricating greases by high-performance liquid chromatography 199
Cantafora, A.
——, Di Biase, A., Alvaro, D. and Angelico, M.
Improved method for measuring the glycine and taurine conjugates of bile salts by high-performance liquid chromatography with tauro-7α,12α-dihydroxy-5β-cholanic acid as internal standard 367
Carpenter, M. W., see O'Laughlin, J. W. 173
Cheung, C.-K., see Pang, C.-P. 309
Chow, J.
——, Orenberg, J. B. and Nugent, K. D.
Comparison of automated pre-column and post-column analysis of amino acid oligomers 243
Cordis, G. A.
——, Engelman, R. M. and Das, D. K.
Simultaneous quantification of myocardial adenine nucleotides and creatinine phosphate by ion-pair reversed-phase high-performance liquid chromatography 283
Das, D. K., see Cordis, G. A. 283
DeGraw, S. R., see Houghten, R. A. 223
Di Biase, A., see Cantafora, A. 367
Dong, S., see Jin, Z. 185
Engelman, R. M., see Cardis, G. A. 283
Eriksson, B.-M.
—— and Person, B.-A.
Liquid chromatographic method for the determination of 3,4-dihydroxyphenylethylene glycol and 3,4-dihydroxymandelic acid in plasma 1
Fetzer, J. C.
—— and Biggs, W. R.
Retention behavior of large polycyclic aromatics in bonded-phase high-performance liquid chromatography 87
——, see Jinno, K. 123
Fiedler, H.-P.
—— and Plaga, A.
Separation of amino acids and antibiotics by narrow-bore and normal-bore high-performance liquid chromatography with pre-column derivatization 229
Field, L. R., see Lee, S. H. 137
Flynn, J. S.
—— and Freeman, D. H.
Mobile phase selectivity mapping of nickel and vanadyl metalloporphyrins in reversed-phase liquid chromatography 111
Freeman, D. H., see Flynn, J. S. 111
Fujie, T., see Abe, Y. 11
Fürst, P., see Brückner, H. 251
Gazdag, M., see Vigh, Gy. 353
Gill, R.
——, Osselton, M. D., Smith, R. M. and Hurdley, T. G.
Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–ammonium nitrate eluent. Interlaboratory collaborative study 65
Gonzalez, M. J.
——, Jaulmes, A., Valentin, P. and Vidal-Madjar, C.
Modelling of peak profiles. Application to the preparative liquid chromatography of steroids 333
Graser, T., see Brückner, H. 251
Grayesky, M. L., see Mann, B. 149
Griest, W. H., see Tomkins, B. A. 103
Guo, D.
——, Mant, C. T. and Hodges, R. S.
Effects of ion-pairing reagents on the prediction of peptide retention in reversed-phase high-performance liquid chromatography 205

Hage, D. S.
—— and Walters, R. R.
Dual-column determination of albumin and immunoglobulin G in serum by high-performance affinity chromatography 37
Ho, K.-C., see Pang, C.-P. 309
Hodges, R. S., see Guo, D. 205
Hoshino, H., see Kanbayashi, M. 191
Houghten, R. A.
—— and DeGraw, S. T.
Effect of positional environmental domains on the variation of high-performance liquid chromatographic peptide retention coefficients 223
Hurdley, T. G., see Gill, R. 65
Ishida, Y., see Ueda, T. 273
Ito, N.
——, Noguchi, K., Kazama, M., Shimura, K. and Kasai, K.-I.
Combination of high-performance affinity chromatography and specific detection of proenzyme applicable to the analysis of the fibrinolytic system of human plasma 51
Jaulmes, A., see Gonzalez, M. J. 333
Jin, Z.
——, Dong, S., Xu, W., Li, Y. and Xu, X.
Analysis of mutagenic nitroarenes in carbon black by high-performance liquid chromatography 185
Jinno, K.
——, Nagoshi, T., Tanaka, N., Okamoto, M., Fetzer, J. C. and Biggs, W. R.
Elution behaviour of peropyrene-type polycyclic aromatic hydrocarbons in various chemically bonded stationary phases in reversed-phase liquid chromatography 123
Jones, G., see Pang, C.-P. 309
Kanbayashi, M.
——, Hoshino, H. and Yotsuyanagi, T.
Highly selective determination of trace amounts of copper(II), nickel(II) and vanadium(V) ions with tetradentate Schiff-base ligands by reversed-phase high-performance liquid chromatography and spectrophotometric detection 191
Kaneko, A., see Abe, Y. 11
Kasai, K.-I., see Ito, N. 51
Kazama, M., see Ito, N. 51
Kharouba, S. N., see Kim, C. 25
Kim, C.
——, Speisky, M. B. and Kharouba, S. N.
Rapid and sensitive method for measuring norepinephrine, dopamine, 5-hydroxytryptamine and their major metabolites in rat brain by high-performance liquid chromatography. Differential effect of probenecid, haloperidol and yohimbine on the concentrations of biogenic amines and metabolites in various regions of rat brain 25
Kobayashi, E., see Abe, Y. 11
Koch, D. D.
—— and Polzin, G. L.
Effect of sample preparation and liquid chromatography column choice on selectivity and precision of plasma catecholamine determination 19
Kříž, J.
——, Vodička, L., Burda, J. and Adamcová, E.
High-performance liquid chromatography of esters of aliphatic and aromatic carboxylic acids 159
Lee, S. H.
—— and Field, L. R.
Fluorescence detection of some nitroamines in high-performance liquid chromatography after post-column reaction 137
Leskinen, L., see Moats, W. A. 79
Li, Y., see Jin, Z. 185
Lim, C. K., see Setchell, K. D. R. 315
Mann, B.
—— and Grayeski, M. L.
New chemiluminescent derivatizing agent for the analysis of aldehydes and ketones by high-performance liquid chromatography with peroxyoxalate chemiluminescence 149
Mant, C. T., see Guo, D. 205
Matsushima, Y., see Abe, Y. 11
Moats, W. A.
—— and Leskinen, L.
Comparison of bonded, polymeric and silica columns for chromatography of some penicillins 79
Nagata, Y., see Abe, Y. 11
Nagoshi, T., see Jinno, K. 123
Nijhof, J., see Wynants, J. 297
Nugent, K. D., see Chow, J. 243
Noguchi, K., see Ito, N. 51
Okamoto, M., see Jinno, K. 123
O'Laughlin, J. W.
—— and Carpenter, M. W.
High-performance liquid chromatographic separation and quantitation of cyanide as the dicyanobis(1,10-phenanthroline)-iron(II) complex 173
Orenberg, J. B., see Chow, J. 243
Osselton, D. M., see Gill, R. 65
Otsubo, O., see Abe, Y. 11
Pang, C.-P.
——, Ho, K.-C., Jones, G. and Cheung, C.-K.
Analysis of total hydroxyproline in urine by high-performance liquid chromatography and pre-column derivatization 309
Perrett, D.
Determination of adenosine ribo- and deoxyribonucleotides as their 1-N^6-etheno derivatives by reversed-phase ion-pair high-performance liquid chromatography 289

Persson, B.-A., see Eriksson, B.-M. 1
Petrov, B., see Wynants, J. 297
Plaga, A., see Fiedler, H.-P. 229
Polzin, G. L., see Koch, D. D. 19
Rasmussen, R. R.
—— and Scherr, M. H.
Preparative low-pressure chromatography of antibiotics on a column of diol-bonded silica gel 325
Rosselli, P., see Valenza, T. 363
Scherr, M. H., see Rasmussen, R. R. 325
Setchell, K. D. R.
——, Welsh, M. B. and Lim, C. K.
High-performance liquid chromatographic analysis of phytoestrogens in soy protein preparations with ultraviolet, electrochemical and thermospray mass spectrometric detection 315
Severin, G.
Monitoring a potential carcinogen in pharmaceutical formulations at the low part per billion level. High-performance liquid chromatographic determination of N-nitrosohexamethyleneimine in tolazamide 57
Shimura, K., see Ito, N. 51
Slégel, P., see Valkó, K. 345
Smith, R. M., see Gill, R. 65
Speisky, M. B., see Kim, C. 25
Suzuki, N., see Abe, Y. 11
Szepesi, G., see Vigh, Gy. 353
Takai, N., see Abe, Y. 11
Tanaka, N., see Jinno, K. 123
Tomkins, B. A.
—— and Griest, W. H.
Liquid chromatographic determination of benzo[*a*]pyrene at part-per-billion concentrations in highly refined coal- and petroleum-derived fuels 103
Ueda, T.
—— and Ishida, Y.
High-performance liquid chromatographic separation of biological macromolecules on new silica-based ion exchangers 273
Valentin, P., see Gonzalez, M. J. 333
Valenza, T.
—— and Rosselli, P.
Rapid and specific high-performance liquid chromatographic determination of clonazepam in plasma 363
Valkó, K.
——, Slégel, P. and Báti, J.
Displacement chromatography of oligomycins 345
Van Belle, H., see Wynants, J. 297
Varga-Puchony, Z., see Vigh, Gy. 353
Vidal-Madjar, C., see Gonzalez, M. J. 333
Vigh, Gy.
——, Varga-Puchony, Z., Szepesi, G. and Gazdag, M.
Semi-preparative high-performance reversed-phase displacement chromatography of insulins 353
Vodička, L., see Kříž, J. 159
Walters, R. R., see Hage, D. S. 37
Watanabe, T., see Abe, Y. 11
Welsh, M. B., see Setchell, K. D. R. 315
Wynants, J.
——, Petrov, B., Nijhof, J. and Van Belle, H.
Optimization of a high-performance liquid chromatographic method for the determination of nucleosides and their catabolites. Application to cat and rabbit heart perfusates 297
Yotsuyanagi, T., see Kanbayashi, M. 191
Xu, W., see Jin, Z. 185
Xu, X., see Jin, Z. 185